

2010 BCBSIL Hospital Profile: Hospital Quality Alliance Measures Benchmarks For Peer Groups 1 - 6 Hospitals - FINAL

**Based on 2010 Annual BCBSIL Hospital Profile
Source: <http://www.hospitalcompare.hhs.gov/>
Current Collection Dates: April 2008-March 2009**

| Hospital Quality Alliance (HQA) Measure | Min Score (%) | State Mean (%) | Median Score (%) | National 90th Percentile (%) | Max Score (%) |
|---|------------------------------------|----------------|------------------|------------------------------|---------------|
| Acute Myocardial Infarction/Heart Attack | | | | | |
| Asprin at Arrival | 79 | 94 | 99 | 100 | 100 |
| Aspirin at Discharge | 70 | 92 | 99 | 100 | 100 |
| Angiotensin Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) for Left Ventricular Systolic Dysfunction (LVSD) | 72 | 89 | 95 | 100 | 100 |
| Beta Blocker at Arrival | No longer part of HQA measure list | | | | |
| Beta Blocker at Discharge | 70 | 93 | 99 | 100 | 100 |
| Percutaneous Coronary Intervention (PCI) Received Within 90 Minutes of Hospital Arrival | 91 | 77 | 93 | 98 | 98 |
| Smoking Cessation Advice/Counseling | 91 | 94 | 100 | 100 | 100 |
| Heart Failure | | | | | |
| Evaluation of Left Ventricular Systolic (LVS) Function | 82 | 93 | 99 | 100 | 100 |
| Angiotensin Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) for Left Ventricular Systolic Dysfunction (LVSD) | 59 | 89 | 95 | 100 | 100 |
| Discharge Instructions | 18 | 79 | 85 | 99 | 100 |
| Smoking Cessation Advice/Counseling | 65 | 93 | 100 | 100 | 100 |
| Pneumonia | | | | | |
| Oxygenation Assessment | No longer part of HQA measure list | | | | |
| Initial Antibiotic Timing | 71 | 94 | 96 | 100 | 100 |
| Pneumococcal Vaccination | 5 | 83 | 89 | 99 | 100 |
| Influenza Vaccination | 3 | 83 | 90 | 99 | 100 |
| Blood Culture Performed in the Emergency Department Prior To Initial Antibiotic Received in Hospital | 68 | 93 | 96 | 100 | 100 |
| Appropriate Initial Antibiotic Selection | 29 | 87 | 89 | 98 | 99 |
| Smoking Cessation Advice/Counseling | 65 | 90 | 97 | 100 | 100 |
| Surgical Infection Prevention | | | | | |
| Preventing Infection | | | | | |
| Prophylactic Antibiotic Received Within 1 Hour Prior to Surgical Incision | 43 | 88 | 95 | 99 | 100 |
| Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time | 54 | 87 | 91 | 99 | 99 |
| Prophylactic Antibiotic Selection | 84 | 94 | 98 | 100 | 100 |
| Cardiac Surgery Patients With Controlled 6 A.M. Postoperative Blood Glucose | 60 | 90 | 92 | 100 | 100 |
| Surgery Patients with Appropriate Hair Removal | 62 | 95 | 100 | 100 | 100 |
| Preventing Blood Clots | | | | | |
| Surgery Patients with Recommended Venous Thromboembolism Prophylaxis Ordered | 4 | 86 | 95 | 99 | 100 |
| Surgery Patients Who Received Appropriate Venous Thromboembolism Prophylaxis Within 24 Hours Prior to Surgery to 24 Hours After Surgery | 5 | 83 | 91 | 98 | 100 |
| Managing Heart Drugs During Surgery | | | | | |
| Surgery Patients on a Beta Blocker Prior to Arrival Who Received a Beta Blocker During the Perioperative Period* | 14 | 86 | 92 | 100 | 100 |

* New measure for 2010 Hospital Profile for information only