

2009 BCBSIL Hospital Profile: Hospital Quality Alliance Measures Benchmarks For Rural Hospitals

Based on 2009 Annual BCBSIL Hospital Profile

Source: <http://www.hospitalcompare.hhs.gov/>

Current Collection Dates: April 2007-March 2008

Hospital Quality Alliance (HQA) Measure	Min Score	State Mean	Median Score	National 90th Percentile	Max Score
Heart Attack					
Heart Attack Patients Given Aspirin at Arrival	86	92	96	100	100
Heart Attack Patients Given Aspirin at Discharge	84	90	97	100	100
Heart Attack Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	94	86	94.5	100	95
Heart Attack Patients Given Beta Blocker at Arrival	76	89	93	100	100
Heart Attack Patients Given Beta Blocker at Discharge	88	93	98	100	100
Heart Attack Patients Given PCI Within 90 Minutes Of Arrival	56	67	62	93	68
Heart Attack Patients Given Smoking Cessation Advice/Counseling	98	90	100	100	100
Heart Failure					
Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	79	87	90	100	100
Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function	17	92	93	100	100
Heart Failure Patients Given Discharge Instructions	44	77	84	97	100
Heart Failure Patients Given Smoking Cessation Advice/Counseling	97	92	100	100	100
Pneumonia					
Pneumonia Patients Given Oxygenation Assessment	67	99	100	100	100
Pneumonia Patients Assessed and Given Pneumococcal Vaccination	34	77	84	97	100
Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior To The Administration Of The First Hospital Dose Of Antibiotics	75	91	95	99	100
Pneumonia Patients Given Smoking Cessation Advice/Counseling	38	88	96	100	100
Pneumonia Patients Given Initial Antibiotic(s) within 4 Hours After Arrival	80	93	96	100	100
Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)	47	87	86	97	100
Pneumonia Patients Assessed and Given Influenza Vaccination	32	76	84	97	100
Surgical Infection Prevention					
Surgery Patients Who Received Preventative Antibiotic(s) One Hour Before Incision	63	86	91	98	100
Surgery Patients Whose Preventative Antibiotic(s) are Stopped Within 24 hours After Surgery	9	81	86	98	98
Surgery Patients Who Received the Appropriate Preventative Antibiotic(s) for Their Surgery	48	93	96	99	100
Surgery Patients Who Received Treatment To Prevent Blood Clots Within 24 Hours Before or After Selected Surgeries to Prevent Blood Clots	51	80	86	96	100
Surgery Patients Whose Doctors Ordered Treatments to Prevent Blood Clots (Venous Thromboembolism) For Certain Types of Surgeries	53	84	89	97	100