

2009 BCBSIL Hospital Profile: Hospital Quality Alliance Measures Benchmarks For Urban Hospitals

Based on 2009 Annual BCBSIL Hospital Profile

Source: <http://www.hospitalcompare.hhs.gov/>

Current Collection Dates: April 2007-March 2008

Hospital Quality Alliance (HQA) Measure	Min Score	State Mean	Median Score	National 90th Percentile	Max Score
Heart Attack					
Heart Attack Patients Given Aspirin at Arrival	81	92	98	100	100
Heart Attack Patients Given Aspirin at Discharge	70	90	98	100	100
Heart Attack Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	65	86	94	100	100
Heart Attack Patients Given Beta Blocker at Arrival	7	89	95	100	100
Heart Attack Patients Given Beta Blocker at Discharge	74	93	98	100	100
Heart Attack Patients Given PCI Within 90 Minutes Of Arrival	40	67	80	93	96
Heart Attack Patients Given Smoking Cessation Advice/Counseling	83	90	99	100	100
Heart Failure					
Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	58	87	93	100	100
Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function	69	92	98	100	100
Heart Failure Patients Given Discharge Instructions	26	77	79	97	100
Heart Failure Patients Given Smoking Cessation Advice/Counseling	30	92	98.5	100	100
Pneumonia					
Pneumonia Patients Given Oxygenation Assessment	93	99	100	100	100
Pneumonia Patients Assessed and Given Pneumococcal Vaccination	2	77	83	97	100
Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior To The Administration Of The First Hospital Dose Of Antibiotics	52	91	94	99	100
Pneumonia Patients Given Smoking Cessation Advice/Counseling	23	88	96	100	100
Pneumonia Patients Given Initial Antibiotic(s) within 4 Hours After Arrival	68	93	95	100	100
Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)	71	87	90	97	98
Pneumonia Patients Assessed and Given Influenza Vaccination	0	76	81	97	100
Surgical Infection Prevention					
Surgery Patients Who Received Preventative Antibiotic(s) One Hour Before Incision	4	86	93	98	100
Surgery Patients Whose Preventative Antibiotic(s) are Stopped Within 24 hours After Surgery	39	81	85	98	98
Surgery Patients Who Received the Appropriate Preventative Antibiotic(s) for Their Surgery	67	93	96	99	100
Surgery Patients Who Received Treatment To Prevent Blood Clots Within 24 Hours Before or After Selected Surgeries to Prevent Blood Clots	4	80	87	96	99
Surgery Patients Whose Doctors Ordered Treatments to Prevent Blood Clots (Venous Thromboembolism) For Certain Types of Surgeries	4	84	91	97	99