



## HEDIS® Measurement Year 2021 PPO Annual Report

Care Provided to BCBSIL PPO Members		BCBSIL PPO HEDIS Rate	Quality Compass National Average (PPOs and EPOs)
<b>Effectiveness of Care: Prevention and Screening</b>			
• Breast Cancer Screening (BCS)		71%	70%
• Cervical Cancer Screening (CCS)		77%	73%
• Colorectal Cancer Screening (COL)		63%	61%
• Chlamydia Screening in Women (CHL) - Total		47%	44%
• Immunizations for Adolescents (IMA) - Combination 2		36%	30%
<b>Effectiveness of Care: Respiratory Conditions</b>			
• Appropriate Testing for Pharyngitis (CWP) - Total		73%	66%
• Pharmacotherapy Management of COPD Exacerbation (PCE)			
○ Systemic Corticosteroid		77%	75%
○ Bronchodilator		83%	81%
• Asthma Medication Ratio (AMR) - Total		87%	83%
<b>Effectiveness of Care: Cardiovascular Conditions</b>			
• Statin Therapy for Patients with Cardiovascular Disease (SPC)			
○ Received Statin Therapy (Total)		85%	82%
○ Statin Adherence 80% (Total)		81%	81%
<b>Effectiveness of Care: Diabetes</b>			
• Comprehensive Diabetes Care (CDC)			
○ Hemoglobin A1c (HbA1c) Testing		89%	89%
• Statin Therapy for Patients with Diabetes (SPD)			
○ Received Statin Therapy		66%	64%
• Statin Adherence 80%		71%	76%
<b>Effectiveness of Care: Behavioral Health</b>			
• Antidepressant Medication Management (AMM)			
○ Effective Continuation Phase Treatment		61%	62%
• Follow-Up Care for Children Prescribed ADHD Medications (ADD)			
○ Continuation & Maintenance Phase		53%	44%
• Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)			
○ Blood Glucose and Cholesterol Testing		36%	35%
• Follow-Up After Hospitalization for Mental Illness - 7 days (Total) (FUH)		58%	48%
• Follow-Up After Emergency Department Visit for Mental Illness - 7 days (Total) (FUM)		56%	48%
• Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence - 7 days (Total) (FUA)		12%	11%
• Follow-Up After High-Intensity Care for Substance Use Disorder - 7 days (Total) (FUI)		43%	47%



## HEDIS® Measurement Year 2021 PPO Annual Report

(Continued from Page 1)

Care Provided to BCBSIL PPO Members		BCBSIL PPO HEDIS Rate	Quality Compass National Average (PPOs and EPOs)
<b>Effectiveness of Care:</b>		<b>Overuse/Appropriateness</b>	
<ul style="list-style-type: none"> <li>Appropriate Treatment for Upper Respiratory Infection (URI) - Total</li> </ul>	89%	87%	
<ul style="list-style-type: none"> <li>Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis (AAB) - Total</li> </ul>	46%	47%	
<ul style="list-style-type: none"> <li>Use of Imaging Studies for Low Back Pain (LBP)</li> </ul>	75%	77%	
<ul style="list-style-type: none"> <li>Avoiding Opioids at High Dosage (HDO)</li> </ul>	97%	95%	
<ul style="list-style-type: none"> <li>Avoiding Opioids from Multiple Prescribers and Multiple Pharmacies (UOP)</li> </ul>	99%	99%	
<ul style="list-style-type: none"> <li>Avoiding potentially risky Continued Opioid Use (COU) - 31 Days (Total)</li> </ul>	98%	98%	
<b>Access/Availability of Care:</b>			
<ul style="list-style-type: none"> <li>Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment (IET)               <ul style="list-style-type: none"> <li>Engagement of AOD - Total</li> </ul> </li> </ul>	16%	13%	
<ul style="list-style-type: none"> <li>Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) - Total</li> </ul>	75%	61%	

Data based on Measurement Year (MY) 2021.

Results are rounded to the nearest percentage.

HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).